

JRSM Short Reports

<http://shr.sagepub.com/>

An evidence-based review of botulinum toxin (Botox) applications in non-cosmetic head and neck conditions

Ricardo Persaud, George Garas, Sanjeev Silva, Constantine Stamatoglou, Paul Chatrath and Kalpesh Patel

JRSM Short Reports 2013 4:

DOI: 10.1177/2042533312472115

The online version of this article can be found at:

<http://shr.sagepub.com/content/4/2/10>

Published by:



<http://www.sagepublications.com>

On behalf of:



[The Royal Society of Medicine](http://www.rsm.ac.uk)

Additional services and information for *JRSM Short Reports* can be found at:

Email Alerts: <http://shr.sagepub.com/cgi/alerts>

Subscriptions: <http://shr.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

>> [Version of Record](#) - Feb 1, 2013

[What is This?](#)



An evidence-based review of botulinum toxin (Botox) applications in non-cosmetic head and neck conditions

Ricardo Persaud¹ • George Garas¹ • Sanjeev Silva² • Constantine Stamatoglou¹ • Paul Chatrath³ • Kalpesh Patel¹

¹Department of Otorhinolaryngology and Head & Neck Surgery, St Mary's Hospital, Imperial College Healthcare NHS Trust, London, UK

²Department of Otorhinolaryngology and Head & Neck Surgery, Manchester Royal Infirmary, Central Manchester University Hospitals NHS Foundation Trust, Manchester, UK

³Department of Otorhinolaryngology and Head & Neck Surgery, Queen's Hospital, Barking, Havering and Redbridge University Hospitals NHS Trust, Essex, UK

Correspondence to: Dr George Garas. Email: georgegaras2@gmail.com

DECLARATIONS

Competing interests

None declared

Funding

None declared

Ethical approval

Not applicable

Guarantor

RP

Contributorship

RP and GG conceived the idea and wrote the initial draft of the paper after reviewing most of the relevant literature. SS, CS and PC also reviewed some of the relevant literature and contributed to the manuscript as well as editing various drafts. KP had access to the data, reviewed and corrected the

Summary

Botulinum toxin (Botox) is an exotoxin produced from *Clostridium botulinum*. It works by blocking the release of acetylcholine from the cholinergic nerve end plates leading to inactivity of the muscles or glands innervated. Botox is best known for its beneficial role in facial aesthetics but recent literature has highlighted its usage in multiple non-cosmetic medical and surgical conditions. This article reviews the current evidence pertaining to Botox use in the head and neck. A literature review was conducted using The Cochrane Controlled Trials Register, Medline and EMBASE databases limited to English Language articles published from 1980 to 2012. The findings suggest that there is level 1 evidence supporting the efficacy of Botox in the treatment of spasmodic dysphonia, essential voice tremor, headache, cervical dystonia, masticatory myalgia, sialorrhoea, temporomandibular joint disorders, bruxism, blepharospasm, hemifacial spasm and rhinitis. For chronic neck pain there is level 1 evidence to show that Botox is ineffective. Level 2 evidence exists for vocal tics, trigeminal neuralgia, dysphagia and post-laryngectomy oesophageal speech. For stuttering, 'first bite syndrome', facial nerve paresis, Frey's syndrome, oromandibular dystonia and palatal/stapedial myoclonus the evidence is level 4. Thus, the literature highlights a therapeutic role for Botox in a wide range of non-cosmetic conditions pertaining to the head and neck (mainly level 1 evidence). With ongoing research, the spectrum of clinical applications and number of people receiving Botox will no doubt increase. Botox appears to justify its title as 'the poison that heals'.

Introduction

Botulinum toxin (Botox) is a protease exotoxin produced from *Clostridium botulinum*. It works by blocking the release of acetylcholine from

cholinergic nerve endings causing inactivity of muscles or glands. Its effects are transient and may be graded by varying the dose and frequency of administration. Botox is one of the most potent naturally occurring biological poisons and in the

manuscript prior to
submission

Reviewers

Nektarios Mazarakis
and Shahab Khan

past has been responsible for many accidental deaths prior to its discovery in medicine. Its first medical use was to treat strabismus in 1980. Nine years later, the cosmetic effects of the toxin on wrinkles were noted, but it was only in 2002, following Food and Drug Administration approval, that Botox gained widespread popularity as an alternative to cosmetic surgery.¹

Recently, the therapeutic uses of Botox have expanded exponentially to include a wide range of medical and surgical conditions. This has been aided by a greater understanding of its underlying physiology as well as improved efficacy and safety. This review examines the evidence on Botox usage in non-cosmetic conditions of the head and neck.

Methods

The Cochrane Controlled Trials Register, Medline and EMBASE databases were searched from 1980 to 2012. The medical subject heading search terms were 'botox' and 'larynx' or 'dysphonia' or 'dystonia' or 'tremor' or 'oral' or 'myoclonus' or 'temporomandibular' or 'sialorrhoea' or 'bruxism' or 'oesophagus' or 'dysphagia' or 'speech' or 'face' or 'autonomic nervous system' or 'sweating' or 'torticollis' or 'pain' or 'migraine' or 'headache' or 'myalgia' or 'neuralgia' or 'nose' or 'rhinitis'. A total of 997 English language abstracts were reviewed and 88 relevant articles identified. Further references were obtained through their bibliographies. Evidence levels, based on those suggested by the Oxford Centre for Evidence-Based Medicine (Table 1),² are shown in the text inside []. The highest level of evidence pertaining to Botox treatment for each of the ENT conditions is presented in Table 2.

Results and discussion

Laryngeal conditions

Spasmodic dysphonia

Spasmodic dysphonia is due to inappropriate glottic closure or opening due to spasm of intrinsic laryngeal muscles. Symptoms include hoarseness and strangled speech breaks (adductor type) or hypophonia and breathy voice (abductor type).³ A meta-analysis of 30 randomized controlled trials (RCTs) involving Botox therapy in adductor spasmodic dysphonia revealed an improvement

to about one standard deviation across the dependent voice-related Quality of Life (QoL) variables studied [1a].^{4,5} A subsequent RCT also confirmed the beneficial effects of Botox in spasmodic dysphonia with the greatest improvements present in those patients who were most profoundly impaired [1b].⁶ In addition, a recent prospective study ($n = 133$) has demonstrated a mean Voice Handicap Index improvement of 9.6% following laryngeal Botox injection in patients with spasmodic dysphonia.⁷

Essential voice tremor

Essential voice tremor is characterized by rhythmic activation of mainly the intrinsic laryngeal muscles. The voice is affected by breaks in pitch, diminished fluency and arrests. It naturally accompanies the ageing process, but may also occur with spasmodic dysphonia.⁸ Electromyography (EMG)-guided Botox injection into the thyroarytenoid muscles was shown to have beneficial effect in a RCT ($n = 13$) [1b],⁹ in a prospective crossover study ($n = 10$) [3b]¹⁰ and a case report [4].¹¹

Stuttering or stammering

This refers to a disorder of speech-motor control in which the flow of speech is disrupted by involuntary repetitions and prolongations of sounds, syllables, words or phrases, with occasional involuntary silent pauses, collectively caused by poor coordination between lingual, labial, laryngeal and respiratory muscles. There is only one case series that has shown that intralaryngeal Botox injection improves fluency in speech therapy failures so its value in treating this disorder is questionable and requires further research [4].¹²

Vocal tics (Gille de la Tourette syndrome)

Repetitive dyskinetic movements of the laryngeal musculature lead to the production of embarrassing speech known as vocal tics. This is commonly seen in Gille de la Tourette syndrome. There is one RCT showing that Botox injections into the thyroarytenoid muscles is efficacious in reducing the frequency and urge of vocal and motor tics ($n = 18$) [2b], but the patients did not report an overall benefit from the treatment.¹³⁻¹⁵ Again, further research is mandated to assess the efficacy of Botox for vocal tics.

Table 1
Levels of evidence based on those suggested by the Oxford Centre for Evidence-Based Medicine

Level of evidence	Type of study
1a	Systematic review (SR) (with homogeneity*) of randomized control trials (RCTs)
1b	Individual RCT (with narrow confidence interval)
1c	All or none [†]
2a	SR (with homogeneity*) of cohort studies
2b	Individual cohort study (including low quality RCT; e.g. <80% follow-up)
2c	'Outcomes' Research; ecological studies
3a	SR (with homogeneity*) of case-control studies
3b	Individual case-control study
4	Case-series, case reports and poor quality cohort or poor quality case-control studies [‡]
5	Expert opinion without explicit critical appraisal, or based on physiology, bench research or 'first principles'

*This refers to a systematic review that is free of worrisome variations (heterogeneity) in the directions and degrees of results between individual studies. Not all systematic reviews with statistically significant heterogeneity need to be worrisome, and not all worrisome heterogeneity need be statistically significant. As noted above, studies displaying worrisome heterogeneity should be tagged with a '-' at the end of their designated level

[†]Met when all patients died before the treatment became available, but some now survive on it; or when some patients died before the treatment became available, but none now die on it

[‡]This refers to a cohort study that failed to clearly define comparison groups and/or failed to measure exposures and outcomes in the same (preferably blinded), objective way in both exposed and non-exposed individuals and/or failed to identify or appropriately control known confounders and/or failed to carry out a sufficiently long and complete follow-up of patients. Poor quality case-control study refers to one that failed to clearly define comparison groups and/or failed to measure exposures and outcomes in the same (preferably blinded), objective way in both cases and controls and/or failed to identify or appropriately control known confounders

Pain

Headache

Numerous multicentre double-blind placebo-controlled trials support the use of Botox as a prophylactic therapy for migraine [1a].¹⁶⁻¹⁸ The technique involves injections into muscles innervated by the facial or trigeminal nerves (e.g. procerus, corrugator, frontalis, temporalis and suboccipital), specific sites of pain distribution or a combination of both.¹⁹ Significant reductions from baseline were observed in patients in the

Botox trial arm with regard to headache and migraine days, cumulative hours of headache and frequency of moderate/severe headache days. A recent meta-analysis confirmed these beneficial effects of Botox but only in the treatment of chronic daily headaches and chronic migraines (>15 episodes per month) [1a]. Adverse effects including blepharoptosis, skin tightness, paraesthesias, neck stiffness, muscle weakness and neck pain can occur at injection sites but these were minimal and transient.²⁰

Cervical dystonia or spasmodic torticollis

This refers to sustained neck muscle contraction resulting in involuntary movements of the head and neck associated with significant cervical pain and abnormal cervical postures. It can be primary or secondary to other neurological disorders.²¹ The evidence supporting the use of Botox in the treatment of cervical dystonia consists of two Cochrane systematic reviews of 13 (677 participants for Botox A) and three (308 participants for Botox B) high-quality RCTs, respectively [1a].^{22,23} These meta-analyses showed that single injection of Botox is effective (as evident from both objective and subjective rating scales) and can be safely repeated if necessary. Since then, there have been further RCTs confirming the efficacy and safety of Botox in the treatment of cervical dystonia in both previously treated as well as Botox-naive patients [1b].²⁴ It is worth noting that Botox not only reduces abnormal movements and contractures but can also prevent secondary degenerative changes of the cervical spine and associated radiculopathy.^{25,26}

Masticatory myalgia

Masticatory pain can be explained by chronic nociceptive irritation of the tendons and fascias of the masseter, temporalis and medial pterygoid muscles.^{27,28} There are three RCTs showing Botox to be more effective than placebo (saline) in reducing masticatory myalgia [1b].²⁹⁻³¹ The most recent of these three RCTs also evaluated with EMG the action potentials of the masseter and temporalis muscles and showed that these decreased by nearly 80% on day 14, and by 25% on day 28 following Botox injection.³⁰ Botox causes a disuse atrophy of the affected muscle which relieves tension, improves aerobic metabolism

Table 2
Levels of evidence for the role of Botox in various head and neck conditions

Condition	Highest level of evidence
Laryngeal	
Spasmodic dysphonia ^{4,5}	1a
Essential voice tremor ⁹	1c
Stuttering ¹²	4
Vocal tics ¹³⁻¹⁵	2b
Pain	
Headache ^{16-18,20}	1a
Cervical dystonia/spasmodic torticollis ^{22,23}	1a
Masticatory myalgia ²⁹⁻³¹	1b
Chronic neck pain ³³ (non-beneficial)	1a
Trigeminal neuralgia ³⁴⁻³⁶	2b
First bite syndrome ^{40,41}	4
Oesophageal	
Oesophageal speech postlaryngectomy ⁴⁴⁻⁴⁷	2c
Dysphagia ⁵¹⁻⁵³	2c
Oral	
Sialorrhoea ⁵⁵⁻⁵⁷	1b
Temporomandibular joint disorders ^{29,31,59}	1b
Bruxism ²⁹	1b
Oromandibular dystonia ^{63,64}	4
Palatal/stapedial myoclonus ^{65,66}	4
Facial	
Blepharospasm ⁶⁹⁻⁷¹	1b
Hemifacial spasm ⁷⁵	1b
Facial nerve paresis ⁷⁶⁻⁷⁸	4
Nasal	
Rhinitis ⁷⁹⁻⁸²	1b
Autonomic	
Frey's syndrome ⁸⁵⁻⁹⁰	4

and enables decompression of afferent nociceptive neurons through reduction of substance P-mediated neurogenic inflammation.^{31,32}

Chronic neck pain (no benefit with Botox)

Several studies have assessed the role of intramuscular Botox injections in chronic neck pain; however, no significant beneficial effect has been demonstrated. A recent Cochrane systematic review of nine trials (503 participants) showed that Botox alone was no better than the placebo (saline) for patients with subacute or chronic neck pain

and concluded that the available evidence does not support the use of Botox either as a monotherapy or in combination with any other treatment in patients with subacute or chronic neck pain [1a].³³

Trigeminal neuralgia

The role of Botox in the treatment of drug-refractory trigeminal neuralgia has been evaluated in three studies ($n = 15$, $n = 12$, $n = 8$, respectively).³⁴⁻³⁶ All three studies (including a low-quality RCT) found Botox to be an effective treatment with the majority of the patients reporting a reduction or even disappearance of the pain [2b].³⁴⁻³⁶ Botox was found to be effective in combination with pharmacotherapy, prior to considering more invasive therapies such as surgery or gamma knife radiosurgery.³⁴ As such, Botox is a particularly valuable treatment for elderly patients and those with adverse anaesthetic comorbidities.^{37,38}

First bite syndrome

This is the development of facial pain after the first bite of each meal and is seen after surgery in the parapharyngeal space, especially deep lobe parotidectomy.³⁹ It is probably due to autonomic dysfunction of salivary myoepithelial cells. Intraparotid Botox injection was found to significantly decrease symptom severity and improve the patients' QoL in a case series of five patients and a case report [4].^{40,41}

Oesophageal conditions

Oesophageal speech post-laryngectomy Tracheoesophageal puncture in laryngectomy patients allows excellent quality speech development in most cases. The procedure involves cricopharyngeal myotomy and valve placement. However, postoperative pharyngo-oesophageal spasm can cause failure of tracheoesophageal speech and dysphagia.⁴² Traditionally, this was treated with dilation of the pharyngo-oesophageal segment (POS), pharyngeal myotomy and/or pharyngeal neurectomy.⁴³ More recently, EMG-guided Botox administration that chemically denervates the cricopharyngeus muscle facilitating tracheoesophageal speech and relieving dysphagia has been reported. There are several

prospective^{44–47} and retrospective outcomes research studies⁴⁸ assessing the efficacy of Botox using both subjective (videotaped recordings) and objective (videostroboscopy) outcome measures [2c]. In corroboration, the largest and most recent prospective study consisting of 34 laryngectomized patients showed Botox therapy to be effective in POS voice restoration, especially when combined with speech therapy [2c].⁴⁴ The effects of Botox were shown to be long-lasting with only one patient needing to be re-injected every three months.⁴⁴ These results are promising but further, higher quality studies are needed to establish the true value of Botox in oesophageal speech post-laryngectomy.

Dysphagia

Incoordination of cricopharyngeal contractions at the initiation of swallowing can result in dysphagia, especially in the elderly population. EMG-guided Botox injections either percutaneously⁴⁹ or endoscopically⁵⁰ to the cricopharyngeus muscle were found to be effective in the treatment of dysphagia in a number prospective and retrospective outcomes research studies [2c].^{51–53} Effective toxin administration can predict a successful surgical outcome following cricopharyngeal myotomy.^{51,54} Again, like with oesophageal speech post-laryngectomy, these results are promising but further, higher quality studies are needed before the true value of Botox in dysphagia is determined.

Oral conditions

Sialorrhoea

Sialorrhoea may occur in neurological and other akinetic disorders such as Parkinson's disease and cerebral palsy. There are several RCTs where the efficacy of Botox injections to the parotid and/or submandibular glands in such patients has been demonstrated [1b].^{55–57} The effects last 3–6 months and can be repeated. Injections can also be used for sialorrhoea caused by salivary fistulas and sialadenitis.⁵⁸

Temporomandibular joint disorders

Spasm of the lateral pterygoid muscles may cause temporomandibular joint (TMJ) disc displacement anteriorly resulting in exquisite pain and clicking.

The evidence supporting the use of Botox in the treatment of such TMJ disorders includes multiple RCTs [1b].^{29,31,59} However, injection of Botox into the lateral pterygoid muscle may cause a 'fixed' smile due to diffusion into the superficial facial muscles.⁶⁰

Bruxism

This is characterized by non-functional contact of the mandibular and maxillary teeth resulting in clenching or tooth grinding due to repetitive, unconscious contraction of the masseter and temporalis muscles.⁶¹ There is one RCT ($n = 30$) which has shown Botox to be efficacious in reducing myofascial pain symptoms in bruxers compared with control patients receiving saline placebo injections²⁹ with a second one currently underway [1b].⁶²

Oromandibular dystonia

This disorder is characterized by involuntary, action-induced, tonic or clonic spasms of the masticatory, lingual and pharyngeal musculature. Symptoms include dysphagia, dysarthria, bruxism and tempomandibular joint subluxation. There are case series and case reports [4] showing favourable effects of Botox injections into the lateral pterygoid, anterior belly of digastric, masseter and temporalis muscles.^{63,64} Thus, further higher quality studies are needed to establish the true role of Botox in the treatment of oromandibular dystonia.

Palatal and stapedius myoclonus

Palatal myoclonus is characterized by involuntary palatal contractions, causing clicking tinnitus due to the action of soft palate muscles on the membranous Eustachian tube. Similarly, stapedius myoclonus can cause clicking tinnitus due to the contractions of the stapedius muscle. There are two case reports, one for each type of myoclonus where the use of Botox has been shown to be beneficial in relieving the patients' symptoms [4]. For palatal myoclonus, Botox was injected in the soft palate under EMG guidance,⁶⁵ while for stapedius myoclonus, Botox was placed trans-tympanically into the middle ear on a piece of gelfoam.⁶⁶ In the latter case, the beneficial effects of Botox lasted for four months.

Facial conditions

Blepharospasm

Involuntary contraction of the eyelid muscles typically occurs bilaterally and in patients over 60 years. The orbicularis oculi muscle is most commonly implicated, but upper facial muscles can also be affected. The therapeutic use of Botox in blepharospasm was first described in 1985⁶⁷ and it has since become the treatment of choice.⁶⁸ There are three RCTs demonstrating the superiority of Botox over placebo [1b].⁶⁹⁻⁷¹ A recent Cochrane systematic review has concluded that doing more RCTs to prove the effectiveness of Botox over the placebo (saline) would be unethical due to the high efficacy and obvious benefits of Botox in treating blepharospasm.⁷²

Hemifacial spasm

This is characterized by unilateral, recurrent, involuntary movements of the muscles innervated by the facial nerve. It is usually due to compression of the facial nerve near its origin by an aberrant branch of the posterior inferior cerebellar artery.⁷³ The first study to assess Botox in hemifacial spasm was in 1986.⁷⁴ Since then, there have been several studies, including one RCT which showed Botox to be an effective and safe treatment.⁷⁵ This RCT involved 11 patients and clearly demonstrated the beneficial effect of the Botox over the placebo [1b].

Facial nerve paresis

Botox may be used to induce therapeutic ptosis, thereby protecting the cornea during the acute phase of facial nerve paresis. This is achieved by transcutaneous injection into Mueller's muscle and the levator palpebrae superioris. There are two case series of therapeutic chemodenervation with Botox of these muscles comprising three and 10 patients, respectively.^{76,77} Both showed that Botox administration is beneficial in preventing damage as well as healing of the cornea [4]. In addition, there is one case series of 30 patients showing Botox to reduce synkinesis in aberrant facial nerve regeneration following facial nerve paresis.⁷⁸ In that study, Botox was injected to several synkinetic muscles of patients with facial nerve paresis and all 30 patients experienced improvement after treatment [4].

Nasal conditions

Rhinitis

In a RCT of 39 patients with allergic rhinitis, Botox therapy provided better symptomatic control than steroid injections into each inferior turbinate, both in terms of the duration and degree of symptoms [1b].⁷⁹ In another RCT of 20 patients with idiopathic (vasomotor) rhinitis, topical application of Botox on a sponge significantly reduced rhinorrhoea compared with placebo (saline) but nasal congestion remained unchanged.⁸⁰ Furthermore, in a study of 38 patients with idiopathic rhinitis, Botox displayed a similar degree and duration of efficacy with regard to hypersecretion symptoms to ipratropium bromide.⁸¹ Middle and inferior turbinate injections of Botox were shown to be a highly effective, safe and simple intervention in a RCT of 30 patients with vasomotor rhinitis [1b].⁸² Hence, the role of Botox seems promising in the treatment of allergic and idiopathic rhinitis though several limiting factors prevent its widespread use. These include the mode of administration which can be associated with the requirement of specialized skills and the potential for significant pain (particularly with injection to the inferior and/or middle turbinates) in addition to its high cost.⁸³

Autonomic conditions

Frey's syndrome

This typically occurs after parotid surgery and is caused by aberrant regeneration of postganglionic parasympathetic fibres innervating sympathetic cholinergic sweat glands. The result is sweating, flushing and piloerection while eating (gustatory sweating).⁸⁴ Several case series have demonstrated the efficacy of Botox in Frey's syndrome [4].⁸⁵⁻⁹⁰ The procedure involves injecting the areas of gustatory sweating identified by an iodine-starch test. Further research is needed to assess the efficacy of Botox as a treatment for Frey's syndrome.

Conclusion

The literature highlights a therapeutic role for Botox in a wide range of non-cosmetic conditions pertaining to Otorhinolaryngology and Head & Neck Surgery. With ongoing research, the spectrum of clinical applications and number of

people receiving Botox will no doubt increase. Botox appears to justify its title as 'the poison that heals'.

References

- Lang A. History and uses of botox (botulinum toxin type a). *Lippincotts Case Manag* 2004;**9**:109–12
- Levels of Evidence published by the Centre for Evidence-Based Medicine, University of Oxford in March 2009. The full reference is available in the following website: <http://www.cebm.net/index.aspx?o=1025>
- Rosenfield DB, Donovan DT, Sulek M, Viswanath NS, Inbody GP, Nudelman HB. Neurologic aspects of spasmodic dysphonia. *J Otolaryngol* 1990;**19**:231–6
- Boutsen F, Cannito MP, Taylor M, Bender B. Botox treatment in adductor spasmodic dysphonia: a meta-analysis. *J Speech Lang Hear Res* 2002;**45**:469–81
- Brazeau GA. Is there time for student intellectual development and scholarly pursuits? *Am J Pharm Educ* 2010;**74**:18
- Cannito MP, Woodson GE, Murry T, Bender B. Perceptual analyses of spasmodic dysphonia before and after treatment. *Arch Otolaryngol Head Neck Surg* 2004;**130**:1393–9
- Novakovic D, Waters HH, D'Elia JB, Blitzer A. Botulinum toxin treatment of adductor spasmodic dysphonia: longitudinal functional outcomes. *Laryngoscope* 2011;**121**:606–12
- Wolraich D, Vasile Marchis-Crisan C, Redding N, Khella SL, Mirza N. Laryngeal tremor: co-occurrence with other movement disorders. *ORL J Otorhinolaryngol Relat Spec* 2010;**72**:291–4
- Adler CH, Bansberg SF, Hentz JG, et al. Botulinum toxin type a for treating voice tremor. *Arch Neurol* 2004;**61**:1416–20
- Warrick P, Dromey C, Irish JC, Durkin L, Pakiam A, Lang A. Botulinum toxin for essential tremor of the voice with multiple anatomical sites of tremor: a crossover design study of unilateral versus bilateral injection. *Laryngoscope* 2000;**110**:1366–74
- Barkmeier-Kraemer J, Lato A, Wiley K. Development of a speech treatment program for a client with essential vocal tremor. *Semin Speech Lang* 2011;**32**:43–57
- Brin MF, Stewart C, Blitzer A, Diamond B. Laryngeal botulinum toxin injections for disabling stuttering in adults. *Neurology* 1994;**44**:2262–6
- Kwak CH, Hanna PA, Jankovic J. Botulinum toxin in the treatment of tics. *Arch Neurol* 2000;**57**:1190–3
- Marras C, Andrews D, Sime E, Lang AE. Botulinum toxin for simple motor tics: a randomized, double-blind, controlled clinical trial. *Neurology* 2001;**56**:605–10
- Porta M, Maggioni G, Ottaviani F, Schindler A. Treatment of phonic tics in patients with Tourette's syndrome using botulinum toxin type a. *Neurol Sci* 2004;**24**:420–3
- Aurora SK, Dodick DW, Turkel CC, et al. Onabotulinumtoxin A for treatment of chronic migraine: results from the double-blind, randomized, placebo-controlled phase of the preempt 1 trial. *Cephalalgia* 2010;**30**:793–803
- Diener HC, Dodick DW, Aurora SK, et al. Onabotulinumtoxin A for treatment of chronic migraine: results from the double-blind, randomized, placebo-controlled phase of the preempt 2 trial. *Cephalalgia* 2010;**30**:804–14
- Dodick DW, Turkel CC, DeGryse RE, et al. Onabotulinumtoxin A for treatment of chronic migraine: pooled results from the double-blind, randomized, placebo-controlled phases of the preempt clinical program. *Headache* 2010;**50**:921–36
- Durham PL, Cady R. Insights into the mechanism of onabotulinumtoxin A in chronic migraine. *Headache* 2011;**51**:1573–7
- Jackson JL, Kuriyama A, Hayashino Y. Botulinum toxin A for prophylactic treatment of migraine and tension headaches in adults: a meta-analysis. *JAMA* 2012;**307**:1736–45
- Velickovic M, Benabou R, Brin MF. Cervical dystonia pathophysiology and treatment options. *Drugs* 2001;**61**:1921–43
- Costa J, Espirito-Santo C, Borges A, et al. Botulinum toxin type A therapy for cervical dystonia. *Cochrane Database Syst Rev* 2005;**1**:CD003633
- Costa J, Espirito-Santo C, Borges A, et al. Botulinum toxin type B for cervical dystonia. *Cochrane Database Syst Rev* 2005;**1**:CD004315
- Comella CL, Jankovic J, Truong DD, Hanschmann A, Grafe S. Efficacy and safety of incobotulinumtoxin A (nt 201, Xeomin(r), botulinum neurotoxin type A, without accessory proteins) in patients with cervical dystonia. *J Neurol Sci* 2011;**308**:103–9
- Jankovic J. Botulinum toxin therapy for cervical dystonia. *Neurotox Res* 2006;**9**:145–8
- Ruiz PJ, Castrillo JC, Burguera JA, et al. Evolution of dose and response to botulinum toxin A in cervical dystonia: a multicenter study. *J Neurol* 2011;**258**:1055–7
- Clark GT. Classification, causation and treatment of masticatory myogenous pain and dysfunction. *Oral Maxillofac Surg Clin North Am* 2008;**20**:145–57, v
- Solberg WK. Temporomandibular disorders: masticatory myalgia and its management. *Br Dent J* 1986;**160**:351–6
- Guarda-Nardini L, Manfredini D, Salamone M, Salmaso L, Tonello S, Ferronato G. Efficacy of botulinum toxin in treating myofascial pain in bruxers: a controlled placebo pilot study. *Cranio* 2008;**26**:126–35
- Kurtoglu C, Gur OH, Kurkcu M, Sertdemir Y, Guler-Uysal F, Uysal H. Effect of botulinum toxin-A in myofascial pain patients with or without functional disc displacement. *J Oral Maxillofac Surg* 2008;**66**:1644–51
- von Lindern JJ, Niederhagen B, Berge S, Appel T. Type A botulinum toxin in the treatment of chronic facial pain associated with masticatory hyperactivity. *J Oral Maxillofac Surg* 2003;**61**:774–8
- Bhagal PS, Hutton A, Monaghan A. A review of the current uses of botox for dentally-related procedures. *Dent Update* 2006;**33**:165–8
- Langevin P, Peloso PM, Lowcock J, et al. Botulinum toxin for subacute/chronic neck pain. *Cochrane Database Syst Rev* 2011;**7**:CD008626
- Bohluli B, Motamedi MH, Bagheri SC, et al. Use of botulinum toxin A for drug-refractory trigeminal neuralgia: preliminary report. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;**111**:47–50
- Turk U, Ilhan S, Alp R, Sur H. Botulinum toxin and intractable trigeminal neuralgia. *Clin Neuropharmacol* 2005;**28**:161–2

- 36 Zuniga C, Diaz S, Piedimonte F, Micheli F. Beneficial effects of botulinum toxin type a in trigeminal neuralgia. *Arq Neuropsiquiatr* 2008;**66**:500–3
- 37 Allam N, Brasil-Neto JP, Brown G, Tomaz C. Injections of botulinum toxin type a produce pain alleviation in intractable trigeminal neuralgia. *Clin J Pain* 2005;**21**:182–4
- 38 Ngeow WC, Nair R. Injection of botulinum toxin type a (botox) into trigger zone of trigeminal neuralgia as a means to control pain. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;**109**:e47–50
- 39 Linkov G, Morris LG, Shah JP, Kraus DH. First bite syndrome: incidence, risk factors, treatment, and outcomes. *Laryngoscope* 2012;**122**:1773–8
- 40 Ali MJ, Orloff LA, Lustig LR, Eisele DW. Botulinum toxin in the treatment of first bite syndrome. *Otolaryngol Head Neck Surg* 2008;**139**:742–3
- 41 Lee BJ, Lee JC, Lee YO, Wang SG, Kim HJ. Novel treatment of first bite syndrome using botulinum toxin type a. *Head Neck* 2009;**31**:989–93
- 42 Chao SS, Graham SM, Hoffman HT. Management of pharyngoesophageal spasm with botox. *Otolaryngol Clin North Am* 2004;**37**:559–66
- 43 Blitzer A, Komisar A, Baredes S, Brin MF, Stewart C. Voice failure after tracheoesophageal puncture: management with botulinum toxin. *Otolaryngol Head Neck Surg* 1995;**113**:668–70
- 44 Bartolomei L, Zambito Marsala S, Pighi GP, et al. Botulinum toxin type a: an effective treatment to restore phonation in laryngectomized patients unable to voice. *Neurol Sci* 2011;**32**:443–7
- 45 Lewin JS, Bishop-Leone JK, Forman AD, Diaz EM Jr. Further experience with botox injection for tracheoesophageal speech failure. *Head Neck* 2001;**23**:456–60
- 46 Meleca RJ, Dworkin JP, Zormeier MM, Simpson ML, Shibuya T, Mathog RH. Videostroboscopy of the pharyngoesophageal segment in laryngectomy patients treated with botulinum toxin. *Otolaryngol Head Neck Surg* 2000;**123**:38–43
- 47 Zormeier MM, Meleca RJ, Simpson ML, et al. Botulinum toxin injection to improve tracheoesophageal speech after total laryngectomy. *Otolaryngol Head Neck Surg* 1999;**120**:314–9
- 48 Hamaker RC, Blom ED. Botulinum neurotoxin for pharyngeal constrictor muscle spasm in tracheoesophageal voice restoration. *Laryngoscope* 2003;**113**:1479–82
- 49 Masiero S, Briani C, Marchese-Ragona R, Giacometti P, Costantini M, Zaninotto G. Successful treatment of long-standing post-stroke dysphagia with botulinum toxin and rehabilitation. *J Rehabil Med* 2006;**38**:201–3
- 50 Parameswaran MS, Soliman AM. Endoscopic botulinum toxin injection for cricopharyngeal dysphagia. *Ann Otol Rhinol Laryngol* 2002;**111**:871–4
- 51 Ahsan SE, Meleca RJ, Dworkin JP. Botulinum toxin injection of the cricopharyngeus muscle for the treatment of dysphagia. *Otolaryngol Head Neck Surg* 2000;**122**:691–5
- 52 Moerman M, Callier Y, Dick C, Vermeersch H. Botulinum toxin for dysphagia due to cricopharyngeal dysfunction. *Eur Arch Otorhinolaryngol* 2002;**259**:1–3
- 53 Murry T, Wasserman T, Carrau RL, Castillo B. Injection of botulinum toxin a for the treatment of dysfunction of the upper esophageal sphincter. *Am J Otolaryngol* 2005;**26**:157–62
- 54 Schneider I, Thumfart WF, Pototschnig C, Eckel HE. Treatment of dysfunction of the cricopharyngeal muscle with botulinum a toxin: introduction of a new, noninvasive method. *Ann Otol Rhinol Laryngol* 1994;**103**:31–5
- 55 Lagalla G, Millevolte M, Capecci M, Provinciali L, Ceravolo MG. Botulinum toxin type a for drooling in Parkinson's disease: a double-blind, randomized, placebo-controlled study. *Mov Disord* 2006;**21**:704–7
- 56 Mancini F, Zangaglia R, Cristina S, et al. Double-blind, placebo-controlled study to evaluate the efficacy and safety of botulinum toxin type a in the treatment of drooling in parkinsonism. *Mov Disord* 2003;**18**:685–8
- 57 Ondo WG, Hunter C, Moore W. A double-blind placebo-controlled trial of botulinum toxin b for sialorrhea in Parkinson's disease. *Neurology* 2004;**62**:37–40
- 58 Ellies M, Gottstein U, Rohrbach-Volland S, Arglebe C, Laskawi R. Reduction of salivary flow with botulinum toxin: extended report on 33 patients with drooling, salivary fistulas, and sialadenitis. *Laryngoscope* 2004;**114**:1856–60
- 59 Dengehem C, Maes JM, Raoul G, Ferri J. [Botulinum toxin a: analgesic treatment for temporomandibular joint disorders]. *Rev Stomatol Chir Maxillofac* 2012;**113**:27–31
- 60 Chikhani L, Dichamp J. [Bruxism, temporo-mandibular dysfunction and botulinum toxin]. *Ann Readapt Med Phys* 2003;**46**:333–7
- 61 Behr M, Hahnel S, Faltermeier A, et al. The two main theories on dental bruxism. *Ann Anat* 2012;**194**:216–9
- 62 Song F, Altman DG, Glenny AM, Deeks JJ. Validity of indirect comparison for estimating efficacy of competing interventions: empirical evidence from published meta-analyses. *BMJ* 2003;**326**:472
- 63 Mendes RA, Upton LG. Management of dystonia of the lateral pterygoid muscle with botulinum toxin a. *Br J Oral Maxillofac Surg* 2009;**47**:481–3
- 64 Moller E, Bakke M, Dalager T, Werdelin LM. Oromandibular dystonia involving the lateral pterygoid muscles: four cases with different complexity. *Mov Disord* 2007;**22**:785–90
- 65 Conill Tobias N, Paula Vernetta CD, Garcia Callejo FJ, Marco Algarrá J. Objective tinnitus from palatal myoclonus. Use of botulinum toxin: a case report. *Acta Otorrinolaryngol Esp* 2012;**63**:391–2
- 66 Liu HB, Fan JP, Lin SZ, Zhao SW, Lin Z. Botox transient treatment of tinnitus due to stapedius myoclonus: case report. *Clin Neurol Neurosurg* 2011;**113**:57–8
- 67 Scott AB, Kennedy RA, Stubbs HA. Botulinum a toxin injection as a treatment for blepharospasm. *Arch Ophthalmol* 1985;**103**:347–50
- 68 Jankovic J. Botulinum a toxin in the treatment of blepharospasm. *Adv Neurol* 1988;**49**:467–72
- 69 Fahn S LT, Moslowitz C, Brin M, Bressman S, Burke R, Scott A. Double blind controlled study of botulinum toxin for blepharospasm. *Neurology* 1985;**35**(Suppl):271–2
- 70 Frueh BR, Nelson CC, Kapustiak JF, Musch DC. The effect of omitting botulinum toxin from the lower eyelid in blepharospasm treatment. *Am J Ophthalmol* 1988;**106**:45–7
- 71 Jankovic J. Blepharospasm and oromandibular-laryngeal-cervical dystonia: a controlled trial of botulinum a toxin therapy. *Adv Neurol* 1988;**50**:583–91

- 72 Costa J, Espirito-Santo C, Borges A, *et al.* Botulinum toxin type a therapy for blepharospasm. *Cochrane Database Syst Rev* 2005;1:CD004900
- 73 Illingworth RD, Porter DG, Jakubowski J. Hemifacial spasm: a prospective long-term follow up of 83 cases treated by microvascular decompression at two neurosurgical centres in the United Kingdom. *J Neurol Neurosurg Psychiatry* 1996;60:72–7
- 74 Elston JS. Botulinum toxin treatment of hemifacial spasm. *J Neurol Neurosurg Psychiatry* 1986;49:827–9
- 75 Yoshimura DM, Aminoff MJ, Tami TA, Scott AB. Treatment of hemifacial spasm with botulinum toxin. *Muscle Nerve* 1992;15:1045–9
- 76 Naik MN, Gangopadhyay N, Fernandes M, Murthy R, Honavar SG. Anterior chemodervation of levator palpebrae superioris with botulinum toxin type-a (botox) to induce temporary ptosis for corneal protection. *Eye (Lond)* 2008;22:1132–6
- 77 Reddy UP, Woodward JA. Abobotulinum toxin a (dysport) and botulinum toxin type a (botox) for purposeful induction of eyelid ptosis. *Ophthalm Plast Reconstr Surg* 2010;26:489–91
- 78 Toffola ED, Furini F, Redaelli C, Prestifilippo E, Bejor M. Evaluation and treatment of synkinesis with botulinum toxin following facial nerve palsy. *Disabil Rehabil* 2010;32:1414–8
- 79 Yang TY, Jung YG, Kim YH, Jang TY. A comparison of the effects of botulinum toxin a and steroid injection on nasal allergy. *Otolaryngol Head Neck Surg* 2008;139:367–71
- 80 Rohrbach S, Junghans K, Kohler S, Laskawi R. Minimally invasive application of botulinum toxin a in patients with idiopathic rhinitis. *Head Face Med* 2009;5:18
- 81 Sapci T, Yazici S, Evcimik MF, *et al.* Investigation of the effects of intranasal botulinum toxin type a and ipratropium bromide nasal spray on nasal hypersecretion in idiopathic rhinitis without eosinophilia. *Rhinology* 2008;46:45–51
- 82 Ozcan C, Vayisoglu Y, Dogu O, Gorur K. The effect of intranasal injection of botulinum toxin a on the symptoms of vasomotor rhinitis. *Am J Otolaryngol* 2006;27:314–8
- 83 Nowak K, Szyfter W. [Application of botulinum toxin a in chronic intrinsic rhinitis]. *Otolaryngol Pol* 2011;65:103–5
- 84 Neumann A, Rosenberger D, Vorsprach O, Dazert S. [The incidence of Frey syndrome following parotidectomy: results of a survey and follow-up]. *HNO* 2011;59:173–8
- 85 Beerens AJ, Snow GB. Botulinum toxin a in the treatment of patients with Frey syndrome. *Br J Surg* 2002;89:116–9
- 86 Cantarella G, Berlusconi A, Mele V, Cogiamanian F, Barbieri S. Treatment of Frey's syndrome with botulinum toxin type b. *Otolaryngol Head Neck Surg* 2010;143:214–8
- 87 de Bree R, Duyndam JE, Kuik DJ, Leemans CR. Repeated botulinum toxin type a injections to treat patients with Frey syndrome. *Arch Otolaryngol Head Neck Surg* 2009;135:287–90
- 88 Drobik C, Laskawi R. Frey's syndrome: treatment with botulinum toxin. *Acta Otolaryngol* 1995;115:459–61
- 89 Pomprasit M, Chintrakarn C. Treatment of Frey's syndrome with botulinum toxin. *J Med Assoc Thai* 2007;90:2397–402
- 90 Steffen A, Rotter N, Konig IR, Wollenberg B. Botulinum toxin for Frey's syndrome: a closer look at different treatment responses. *J Laryngol Otol* 2012;126:185–9

© 2013 The Author(s)

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-commercial License (<http://creativecommons.org/licenses/by-nc/2.0/>), which permits non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.